


Substitute for form 1449B/PTO INFORMATION DISCLOSURE STATEMENT BY APPLICANT (use as many sheets as necessary)		<i>Complete if Known</i>			
		Application Number	10/759,519		
		Filing Date	January 16, 2004		
		First Named Inventor	Cantor et al.		
		Group Art Unit	1645		
		Examiner Name	To be assigned		
Sheet		of		Attorney Docket Number	701586-053651

OTHER PRIOR ART - NON PATENT LITERATURE DOCUMENTS			
Examiner Initials*	Cite No. ¹	Include name of the author (in CAPITAL LETTERS), title of the article (when appropriate), title of the item (book, magazine, journal, serial, symposium, catalog, etc.), date, page(s), volume-issue number(s), publisher, city and/or country where published.	T ²
STK	C1	Grupe, A. et al. In silico mapping of complex disease-related traits in mice. <i>Science</i> 292, 1915-8. (2001).	
	C2	Hirschhorn, J.N., Lohmueller, K., Byrne, E. & Hirschhorn, K. A comprehensive review of genetic association studies. <i>Genet Med</i> 4, 45-61. (2002).	
	C3	Zhang, S., Pakstis, A.J., Kidd, K.K. & Zhao, H. Comparisons of two methods for haplotype reconstruction and haplotype frequency estimation from population data. <i>Am J Hum Genet</i> 69, 906-14. (2001).	
	C4	Templeton, A.R., Sing, C.F., Kessling, A. & Humphries, S. A cladistic analysis of phenotype associations with haplotypes inferred from restriction endonuclease mapping. II. The analysis of natural populations. <i>Genetics</i> 120, 1145-54. (1988).	
	C5	Kruglyak, L. Prospects for whole-genome linkage disequilibrium mapping of common disease genes. <i>Nat Genet</i> 22, 139-44. (1999).	
	C6	Judson, R., Stephens, J.C. & Windemuth, A. The predictive power of haplotypes in clinical response. <i>Pharmacogenomics</i> 1, 15-26. (2000).	
	C7	Martin, E.R. et al. Analysis of association at single nucleotide polymorphisms in the APOE region. <i>Genomics</i> 63, 7-12. (2000).	
	C8	Clark, A.G. Inference of haplotypes from PCR-amplified samples of diploid populations. <i>Mol Biol Evol</i> 7, 111-22. (1990).	
	C9	Stephens, M., Smith, N.J. & Donnelly, P. A new statistical method for haplotype reconstruction from population data. <i>Am J Hum Genet</i> 68, 978-89. (2001).	

57K	C10	Ruano, G. & Kidd, K.K. Direct haplotyping of chromosomal segments from multiple heterozygotes via allele-specific PCR amplification. <i>Nucleic Acids Res</i> 17, 8392. (1989).	
	C11	Ruano, G., Kidd, K.K. & Stephens, J.C. Haplotype of multiple polymorphisms resolved by enzymatic amplification of single DNA molecules. <i>Proc Natl Acad Sci U S A</i> 87, 6296-300. (1990).	
	C12	Douglas, J.A., Boehnke, M., Gillanders, E., Trent, J.M. & Gruber, S.B. Experimentally-derived haplotypes substantially increase the efficiency of linkage disequilibrium studies. <i>Nat Genet</i> 28, 361-4. (2001).	
	C13	Stephens, J.C., Rogers, J. & Ruano, G. Theoretical underpinning of the single-molecule-dilution (SMD) method of direct haplotype resolution. <i>Am J Hum Genet</i> 46, 1149-55. (1990).	
	C14	Daly, M.J., Rioux, J.D., Schaffner, S.F., Hudson, T.J. & Lander, E.S. High-resolution haplotype structure in the human genome. <i>Nat Genet</i> 29, 229-32. (2001).	
	C15	Gabriel, S.B. et al. The structure of haplotype blocks in the human genome. <i>Science</i> 296, 2225-9. (2002).	
	C16	Shields and Harris (2000) <i>J. Clin. Onc.</i> 18:2309-2316.	
	C17	Evans and Relling (1999) <i>Science</i> 286:487-491	
	C18	Collins et al. (1997) <i>Science</i> 278, 1580-81	
	C19	Drysdale et al. (2000) <i>Proc. Natl. Acad. Sci.</i> 97:10483-8 / Krynetski and Evans (1998) <i>Am. J. Hum. Gen.</i> 63:11-16).	
	C 20	Orita et al. (1989) <i>Proc. Natl. Acad. Sci. USA</i> 86:2766-2770	
	C21	Prior et al. (1995) <i>Hum. Mutat.</i> 5:263-268	
	C22	Nickerson et al. (1990) <i>Proc. Natl. Acad. Sci. USA</i> 87:8923-892	

511	C23	Conner et al. (1983) Proc. Natl. Acad. Sci. USA 80:278-282	
	C24	Ruano et al. (1990) Proc. Natl. Acad. Sci. USA 87:6296-6300 C11	
	C25	Michalatos-Beloin et al. (1996) Nucleic Acids Res. 24:4841-4843	
	C26	Ruano and Kidd (1991) Nucleic Acids Res. 19:6877-6882	
	C27	Cheng, et al. (1994) Proc. Natl. Acad. Sci. USA 91:5695-5699 ✓	
	C28	Papadopoulos et al. (1995) Nature Genet. 11:99-102	
	C29	Woolley et al. (2000) Nature Biotech. 18:760-763	


Examiner Signature		Date Considered	3-15-06
-----------------------	---	--------------------	---------

* EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

¹ Applicant's unique citation designation number (optional). ² Applicant is to place a check mark here if English language Translation is attached.



FORM PTO-1449				DOCKET NO.: 701586-053651		SERIAL NO.: 10/759,519	
INFORMATION DISCLOSURE STATEMENT				APPLICANT(S): Charles R. Cantor and Chunming Ding			
				FILING DATE: January 16, 2004		GROUP NO.: 1645	
UNITED STATES PATENT DOCUMENTS							
EXAM. INITIALS		DOCUMENT NUMBER	DATE	INVENTOR/ASSIGNEE	CLASS	SUBCLASS	FILING DATE IF APPROPRIATE
FOREIGN PATENT DOCUMENTS							
		DOCUMENT NUMBER	DATE	COUNTRY	CLASS	SUBCLASS	TRANSLATION YES/NO
OTHER DOCUMENTS (INCLUDING AUTHOR, TITLE, DATE, PERTINENT PAGES, ETC.)							
STK	C1	Barnes (1994) Proc. Natl. Acad. Sci. USA 91:2216-2220					
	C2	Clark, A.G. Inference of haplotypes from PCR-amplified samples of diploid populations. Mol Biol Evol 7, 111-22. (1990).					
	C3	Collins et al. (1997) Science 278, 1580-81					
	C4	Conner et al. (1983) Proc. Natl. Acad. Sci. USA 80:278-282					
	C5	Daly, M.J., Rioux, J.D., Schaffner, S.F., Hudson, T.J. & Lander, E.S. High-resolution haplotype structure in the human genome. Nat Genet 29, 229-32. (2001).					
	C6	Douglas, J.A., Boehnke, M., Gillanders, E., Trent, J.M. & Gruber, S.B. Experimentally derived haplotypes substantially increase the efficiency of linkage disequilibrium studies. Nat Genet 28, 361-4. (2001).					
	C7	Drysdale et al. (2000) Proc. Natl. Acad. Sci. 97:10483-8					
	C8	Evans and Relling (1999) Science 286:487-491					
	C9	Gabriel, S.B. et al. The structure of haplotype blocks in the human genome. Science 296, 2225-9. (2002).					
	C10	Grupe, A. et al. In silico mapping of complex disease-related traits in mice. Science 292, 1915-8. (2001).					
	C11	Hirschhorn, J.N., Lohmueller, K., Byrne, E. & Hirschhorn, K. A comprehensive review of genetic association studies. Genet Med 4, 45-61. (2002).					
	C12	Hodge et al. (1999) Nature Genet. 21:360-361					
	C13	Judson, R., Stephens, J.C. & Windemuth, A. The predictive power of haplotypes in clinical response. Pharmacogenomics 1, 15-26. (2000).					



57K		Kruglyak, L. Prospects for whole-genome linkage disequilibrium mapping of common disease genes. Nat Genet 22, 139-44. (1999).
	C15	Krynetski and Evans (1998) Am. J. Hum. Gen. 63:11-16
	C16	Martin, E.R. et al. Analysis of association at single-nucleotide polymorphisms in the APOE region. Genomics 63, 7-12. (2000).
	C17	Michalatos-Beloin et al. (1996) Nucleic Acids Res. 24:4841-4843
	C18	Nickerson et al. (1990) Proc. Natl. Acad. Sci. USA 87:8923-8927
	C19	Orita et al. (1989) Proc. Natl. Acad. Sci. USA 86:2766-2770
	C20	Papadopoulos et al. (1995) Nature Genet. 11:99-102
	C21	Prior et al. (1995) Hum. Mutat. 5:263-268
	C22	Ruano and Kidd (1989) Nucleic Acids Res. 17:8392
	C23	Ruano and Kidd (1991) Nucleic Acids Res. 19:6877-6882
	C24	Ruano et al. (1990) Proc. Natl. Acad. Sci. USA 87:6296-6300
	C25	Shields and Harris (2000) J. Clin. Onc. 18:2309-2316
	C26	Stephens, J.C., Rogers, J. & Ruano, G. Theoretical underpinning of the single-molecule dilution (SMD) method of direct haplotype resolution. Am J Hum Genet 46, 1149-55. (1990).
	C27	Stephens, M., Smith, N.J. & Donnelly, P. A new statistical method for haplotype reconstruction from population data. Am J Hum Genet 68, 978-89. (2001).
	C28	Templeton, A.R., Sing, C.F., Kessling, A. & Humphries, S. A cladistic analysis of phenotype-associations with haplotypes inferred from restriction endonuclease mapping. II. The analysis of natural populations. Genetics 120, 1145-54. (1988).
	C29	Woolley et al. (2000) Nature Biotech. 18:760-763
	C30	Zhang, S., Pakstis, A.J., Kidd, K.K. & Zhao, H. Comparisons of two methods for haplotype reconstruction and haplotype frequency estimation from population data. Am J Hum Genet 69, 906-14. (2001).